

Collection No. 90

Riverina, N.S.W.

K. H. L. Key

3-6/9/59

3/9/59.

57379 Prod. on plain, brown-gray soil,  
<sup>Lupinus sp.</sup>  
puff-shelf-depression complex, widely  
spaced myall, various grasses to 9 in.  
Chort. (n)

389. Narrandera

4/9. 389. Narrandera.

413. Flat plain with brown loam. Trellis,  
<sup>seeds, nov. 22 (?) = p. (j.)</sup>  
prob. originally myall. *Stipa aristiglumis*  
<sup>creeping</sup>  
to 1 ft, acc. of *Atriplex* & *Bassia*. Chort (n).

422. Plain with brown loam. Turned  
<sup>*Cryptobothrus chrysophorus*</sup>  
from licolot, rosewood, pine, ~~to~~  
bullock. *Bassia quinqueangula*, *Stipa*,  
*Chloris acicularis*, barley grass. Oedolus  
(n), Chort (n)

435. Tall *canadensis* lining  
river with muddy banks.

465. Turned off to Currathool.

470. Currathool

475. Back on Short Highway

482. Flat treless plain, brown to grey-brown  
<sup>*Percybrannan*</sup>  
greasy to sm. soil with a dead grass

in tussocks to 6 in. <sup>scattered</sup> *Kochia* sp. to 1 ft., very low green pick of forbs.

492. Flat featureless plain with grey-brown <sup>*Euphorbia* sp. *Eumys* nov. 22(?) sp.</sup>

compact <sup>vol. sm soil</sup> patches of a *Helichrysum apiculatum* (Vahl.) DC. <sup>sens. lat.</sup>

moody composition to 1 ft., a small *Helipterium corymbiflorum* Schlecht.

siliques *Helipterum* (spec.) to 3 in in

between, with occ. low *Bassia* & ~~other~~

tussocks of dead grass to 6 in. *Periclyptera*

*rana* (n) *Morabaskia* on & between the

shrubs, but probably mainly between

them, possibly feeding on the low

<sup>observed feeding on flowers of the shrubby, composite</sup> *Helipterum*. (18 mile peg from Hay)

511. Hay.

519. 512 Hay.

514. Large flowered woodland to soft,

*Periclyptera rana*, with dense *Bassia quinqueangulata* to

18 in, *Atriplex* spp. to 9 in, occ. soft

*bassias*, barley grass. Grey self-mulching

soil with large cracks & holes.

*Schinus* (n) *Chort* (n)

528. Perfectly flat plain, slightly to mod.

<sup>*Eumys* nov. 22(?) sp., *Eumys* nov. 23(?) sp.</sup>

sm grey-brown soil, with dense

patches of

*Kochia, aphylla* & *Atriplex vesicaria* to  
18 in., low annual *Atriplex* spp. & silvery  
*Helipterum*, occ. barley grass. Very dry.

Perelybrana.

539. Flat plain, compact brown loam.

*Perelybrana rana* (f.)

*Bassia* spp. & annual *Atriplex*.

553. Red-brown sandy loam plain.  
*Lycopers* sp. *Ixiolaena tomentosa* Soud. & F. Muell.

with yellow-flowered composite to 10 DC  
to 4 in (sp.) *Helipterum floribundum*  
& white everlasting & a tussocky grass

to 6 in. Perelybrana (v).

562 Gumbao

583. Goolgowi

*Krugeria marida*, *Azylota ochracea* (f.), *Cratilopus*

591. Mallee & stunted *intersecta* to 20 ft  
sp. *Micocela* sp. (to 6 ft) *Gum* not (to 1 ft) 48 sp. 2

with broom bush & *Trochala*, & a very  
& a microphyllous shrub (thrust to 3 ft.)  
narrow-phyllode wattle to 3 ft, Very

sparse grass. Red sandy loam, undulating.

605. *Dealbata*, *Callitris*  
(all to 30 ft) *Poppyifolia* to 10 ft

*glauca* & a phyllode wattle (spike type),  
*Moraba* sp. (P48) *Gum* not. 220 sp.

on rocky quartzitic slope (The Gap?) in

a gap in range. A strongly tussocky grass

(Mt. Boppy species), *san Anotida* sp. Brown

sandy loam. *Moraba* mainly on

tussocky grass.

619. Hauke's Springs.

619. 678. W. Wyalong.

693. *Microcampa* to 40 ft., *Uelab* to 30 ft.,  
*Monistria pustulifera*  
yellow to 10 ft. on flat brown loams  
with *Chloris acicularis* & occ. forbs.

*Monistria* associated with ~~abundant~~  
~~leafed plant (spec.)~~ low growth of  
*Echium plantagineum*.

698 Barmudman.

699. Barmudman cemetery. *Microcampa*,  
*C. glauca*, *Uelab*, *Hypposium* sp.,

*Lupatrine*, *Chloris acicularis*, *Shipa* sp.

No *Helichrysum*, no *Themeda*, no *Moraba*.

722. Timora.

723. Timora cemetery. no *Themeda* or  
*Helichrysum*: not examined for *Moraba*.

757. Stockinbingal Cemetery. *Meliodora* &  
*albens*. Good *Themeda* & *Helichrysum*:  
no *Moraba*.